

The Neutrino Suppression Scale from Large Volumes

Joseph P. Conlon and Daniel Cremades

DAMTP, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA, UK

(Dated: February 2, 2008)

We present an argument in which the scale ~ 0.1 eV associated with neutrino masses naturally appears in a class of (very) large volume compactifications, being tied to a supersymmetry scale of 10^3 GeV and a string scale of 10^{11} GeV. The masses are of Majorana type and there is no right-handed neutrino within the low-energy field theory. The suppression scale 10^{14} GeV is independent of the masses of the heavy states that are integrated out. These kind of constructions appear naturally in Type IIB flux compactifications. However, the arguments that lead to this result rely only on a few geometrical features of the compactification manifold, and hence can be used independently of string theory.

PACS numbers: 11.25.Mj 11.25.Wx 14.60.Pq 14.60.St

I. INTRODUCTION

The existence of non-zero neutrino masses established in recent years constitutes the first evidence of physics beyond the Standard Model (for a review, see [1]). Neutrino mass splittings have been measured [2]

$$\begin{aligned} 7.7 \times 10^{-5} \text{ eV}^2 &\lesssim \Delta m_{12}^2 \lesssim 8.4 \times 10^{-5} \text{ eV}^2, \\ 1.9 \times 10^{-3} \text{ eV}^2 &\lesssim \Delta m_{23}^2 \lesssim 3.0 \times 10^{-3} \text{ eV}^2. \end{aligned}$$

The cosmological constraint $\sum_\nu m_\nu \lesssim 0.7 \text{ eV}$ thus implies the heaviest neutrino has a mass

$$0.05 \text{ eV} \lesssim m_\nu \lesssim 0.3 \text{ eV}. \quad (1)$$

If this mass is of Dirac origin, the largest possible neutrino Yukawa coupling is at least six orders of magnitude smaller than that of the electron. This may be achieved in warped compactifications [3, 4, 5]. However arguably the smallness of neutrino masses points to a Majorana nature, as Majorana neutrinos require the existence of highly suppressed non-renormalisable operators. This also implies the existence of a new physical scale between the electroweak and Planck scales.

In the Standard Model, neutrino masses are generated after electroweak symmetry breaking by the dimension five operator,

$$\mathcal{O}_5 = \frac{\lambda}{\Lambda} (L^T C i \tau_2 \tau L) (H^T i \tau_2 \tau H). \quad (2)$$

In the MSSM, the comparable superpotential operator is (suppressing gauge theory indices)

$$\mathcal{O} = \frac{\lambda}{\Lambda} H_2 H_L L L. \quad (3)$$

These are also the effective operators that appear in the seesaw mechanism once the right-handed neutrinos are integrated out. In the seesaw mechanism Λ is the scale of right-handed neutrino masses where new physics occurs.

The observed scale of neutrino masses creates a puzzle. The MSSM Higgs vev is $\langle H_2 \rangle = v \sin \beta / \sqrt{2}$, with $v =$

246 GeV, which generates neutrino masses

$$m_\nu = \frac{\lambda v^2 \sin^2 \beta}{2\Lambda} = \lambda \sin^2 \beta (0.10 \text{ eV}) \left(\frac{3 \times 10^{14} \text{ GeV}}{\Lambda} \right) \quad (4)$$

If $\sin^2 \beta \sim 1$ (as applicable in the large $\tan \beta$ regime), $m_\nu \sim 0.1 \text{ eV}$ requires a suppression scale $\Lambda = 3 \times 10^{14} \text{ GeV}$ for $\lambda = 1$ (roughly the top Yukawa coupling) and $\Lambda \sim 3 \times 10^{12} \text{ GeV}$ for $\lambda = 0.01$ (roughly the tau Yukawa coupling).

The value of Λ is mysterious as it is neither the GUT scale ($\sim 2 \times 10^{16} \text{ GeV}$), for which neutrino masses are too small, nor the intermediate scale 10^{11} GeV , for which neutrino masses are too large: the scale $10^{12} \text{ GeV} \div 10^{14} \text{ GeV}$ does not correspond to any other physical scale.

In this note we show that in a certain class of higher dimensional constructions, and assuming only that lepton number is not a fundamental symmetry of the low-energy theory, the required scale of Λ emerges naturally from requiring TeV-scale supersymmetry. This class of constructions appears naturally in the context of flux compactifications of Type IIB string theory [6], but the arguments leading to the correct scale of neutrino masses need not be particular to them and rely only on a few geometric features of the model. Thus any construction, stringy or not, that shares these characteristics will give this scale.

These constructions are characterised by a stabilised large compactification manifold, with $\mathcal{V} \sim 10^{15}$ in string units, with an intermediate string scale $m_s \sim 10^{11} \text{ GeV}$. In this respect they have similarities to the large extra dimensions scenario [7, 8] (see [9, 10] for neutrino masses in this scenario and [11, 12] for studies of an intermediate string scale). The Standard Model is localised on branes wrapping small cycles within the large overall volume. Considered in string theory, these constructions present many appealing phenomenological features [6, 13], such as a naturally occurring hierarchy and computable flavour-universal soft terms [14, 15]. The large volume also allows a considerable degree of control and simplification in the computations.

The structure of this note is as follows. In section II we review large volume models as presented in [6], and estimate the strength of the $HHLL$ operator using the

results of [14]. In section III we show that the correct dimensionful suppression scale arises independently of the masses of the heavy fields that are integrated out. In section IV we comment on embedding into a fully fledged string model. In section V we conclude.

II. LARGE VOLUME MODELS AND NEUTRINO MASSES

The large volume models arise in IIB flux compactifications [16] with α' corrections [17] and nonperturbative superpotential terms [18]. The relevant Kähler potential and superpotential are [24]

$$\begin{aligned}\mathcal{K} &= -2 \ln \left(\mathcal{V} + \frac{\xi}{g_s^{3/2}} \right) - \ln \left(i \int \Omega \wedge \bar{\Omega} \right) - \ln(S + \bar{S}), \\ W &= \int G_3 \wedge \Omega + \sum A_i \exp(-a_i T_i).\end{aligned}\quad (5)$$

For the simplest model (the Calabi-Yau $\mathbb{P}^4_{[1,1,1,6,9]}$), there are two Kähler moduli and the volume can be written as

$$\mathcal{V} = \frac{1}{9\sqrt{2}} \left(\left(\frac{T_b + \bar{T}_b}{2} \right)^{3/2} - \left(\frac{T_s + \bar{T}_s}{2} \right)^{3/2} \right). \quad (6)$$

Moduli stabilisation in this geometry has been studied in detail in [6, 13]. The dilaton and complex structure moduli are flux-stabilised. Denoting $\tau_s = \text{Re}(T_s)$, $\tau_b = \text{Re}(T_b)$, the scalar potential for the Kähler moduli is

$$V = \frac{\sqrt{\tau_s} a_s |A_s|^2 e^{-2a_s \tau_s}}{\mathcal{V}} - \frac{a_s \tau_s |A_s W_0| e^{-a_s \tau_s}}{\mathcal{V}^2} + \frac{\xi |W_0|^2}{g_s^{3/2} \mathcal{V}^3}. \quad (7)$$

In this expression, $W_0 = \langle \int G_3 \wedge \Omega \rangle$ and typically is $\mathcal{O}(1)$. The minimum of (7) lies at exponentially large volume,

$$\mathcal{V} \sim W_0 e^{c/g_s}, \quad (8)$$

with $\tau_s \sim \ln \mathcal{V}$ and c an $\mathcal{O}(1)$ constant. The Standard Model lives on D7 branes wrapping the small cycle τ_s .

The gravitino mass is given by

$$m_{3/2} = e^{K/2} W_0 = \frac{W_0}{\mathcal{V}}. \quad (9)$$

A TeV-scale gravitino mass, required for a natural solution to the weak hierarchy problem, requires a volume $\mathcal{V} \sim 10^{15}$. This occurs for appropriate values of g_s and ξ .

The extra-dimensional geometry is then a small four-dimensional region containing the Standard Model, embedded in a large six dimensional Calabi-Yau. We emphasize that while a natural stringy embedding exists, our results will apply to any model giving this geometry.

The value of Λ in the field theory equation (3) depends on the underlying supergravity structure. If we use C to denote generic matter fields (and their scalar

components), on general grounds we can expand the superpotential as a power series in C ,

$$W = \hat{W} + \frac{Y_{\alpha\beta\gamma}}{6} C^\alpha C^\beta C^\gamma + \frac{Z_{\alpha\beta\gamma\delta}}{24 M_P} C^\alpha C^\beta C^\gamma C^\delta + \dots \quad (10)$$

The dimensionful suppression scale in the superpotential must be M_P , as this is the only scale known in supergravity. In stringy constructions this is directly required by holomorphy: other scales such as the string or Kaluza-Klein scales are related to M_P by non-holomorphic functions of the moduli: e.g. $M_s \sim M_P/\mathcal{V}^{1/2} \sim M_P/(T_b + \bar{T}_b)^{2/3}$.

Fermion bilinears arise via the Lagrangian term

$$\int d^4x e^{\hat{K}/2} \partial_\alpha \partial_\beta W \psi^\alpha \psi^\beta. \quad (11)$$

This generates terms

$$e^{\hat{K}/2} Y_{\alpha\beta\gamma} C^\alpha \psi^\beta \psi^\gamma + e^{\hat{K}/2} \frac{Z_{\alpha\beta\gamma\delta}}{2 M_P} C^\alpha C^\beta \psi^\gamma \psi^\delta \in \mathcal{L} \quad (12)$$

The operator $\mathcal{O}$ thus arises from the superpotential term

$$\frac{\lambda}{M_P} H_2 H_2 L L \in W.$$

The main assumption we make is that, consistent with the symmetries of the R-parity MSSM, this operator exists in the superpotential.

Equation (12) does not itself give the physical coupling strength as in general the matter field metrics fail to be canonically normalised. The full Lagrangian is

$$\begin{aligned}\mathcal{L} &= \tilde{K}_{\alpha\beta} \partial_\mu C^\alpha \partial^\mu \bar{C}^\beta + e^{\hat{K}/2} Y_{\alpha\beta\gamma} C^\alpha \psi^\beta \psi^\gamma \\ &+ e^{\hat{K}/2} \frac{Z_{\alpha\beta\gamma\delta}}{M_P} C^\alpha C^\beta \psi^\gamma \psi^\delta.\end{aligned}\quad (13)$$

The physical strengths of the dimension-four (i.e. Yukawa) and dimension five couplings are then

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{1/2}}, \quad (14)$$

$$\hat{Z}_{\alpha\beta\gamma\delta} = e^{\hat{K}/2} \frac{Z_{\alpha\beta\gamma\delta}}{M_P (\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma \tilde{K}_\delta)^{1/2}}. \quad (15)$$

We have restricted (13) to diagonal metrics, but this is only for convenience. The Kähler potentials of both moduli and matter fields are crucial for computing the physical suppression scale of dimension-five operators. In IIB string compactifications, the moduli Kähler potential is

$$\hat{K} = -2 \ln \mathcal{V} - \ln \left(i \int \Omega \wedge \bar{\Omega} \right) - \ln(S + \bar{S}). \quad (16)$$

The factor $-2 \ln \mathcal{V}$ is generic as it ensures the ‘natural’ scale of the scalar potential is m_s^4 rather than M_P^4 (using $m_s \sim \frac{M_P}{\sqrt{\mathcal{V}}}$). It will therefore be present in any model fitting into the framework described in section II.

It was shown in [14] that for the models described in section II, the matter metric scales at leading order as

$$\tilde{K}_{\alpha\bar{\beta}} \sim \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}} k_{\alpha\bar{\beta}}(\phi), \quad (17)$$

where ϕ are complex structure moduli. The volume scaling is easiest to understand. As the Standard Model interactions are localised on a small cycle, the physical Yukawas must be insensitive to the bulk volume. Using (14) and (16), this can hold only if the matter metrics scale as $\tilde{K}_\alpha \sim \mathcal{V}^{-2/3}$ [25]. This is an argument from locality, and so is not specific to string theory and will hold for any model giving this geometry. The power of τ_s depends on the brane configuration - $1/3$ is for the simplest case where all branes wrap the same cycle.

Equation (17) suffices to determine the suppression scale for dimension-five operators. From (15) we find

$$\hat{Z}_{\alpha\beta\gamma\delta} \sim \frac{\mathcal{V}^{1/3}}{M_P \tau_s^{2/3}} Z_{\alpha\beta\gamma\delta}. \quad (18)$$

As in equation (9), we require $\mathcal{V} \sim 10^{15}$ for a natural solution to the weak hierarchy problem. The size of the small cycle τ_s determines the Standard Model gauge couplings by $\alpha_{SM}(m_s) \sim \tau_s^{-1}$ and so we require $\tau_s \sim 20$. The dimension-five operator $\hat{Z}$ is then

$$\begin{aligned} \hat{Z}_{\alpha\beta\gamma\delta} &= \frac{Z_{\alpha\beta\gamma\delta}}{1.7 \times 10^{14} \text{GeV}} \left(\frac{\mathcal{V}}{10^{15}} \right)^{1/3} \left(\frac{20}{\tau_s} \right)^{2/3} \\ \Lambda &= (1.7 \times 10^{14} \text{GeV}) \left(\frac{10^{15}}{\mathcal{V}} \right)^{1/3} \left(\frac{\tau_s}{20} \right)^{2/3} \end{aligned} \quad (19)$$

The operator $\hat{Z}_{\alpha\beta\gamma\delta}$ is defined at the string scale (10^{11}GeV) and renormalised to the neutrino mass scale (10^{-1}eV). MSSM loop corrections involve weakly coupled gauge groups, and so while may alter the detailed form of $\hat{Z}_{\alpha\beta\gamma\delta}$ may alter, the overall scale of Λ should not change. This is analogous to the running of slepton or sneutrino masses, which also get masses from non-renormalisable operators: the physical and high-scale values differ by $\mathcal{O}(1)$ factors. Such a factor does not affect the overall scale of (19). If there exist extra strongly coupled hidden sectors, (19) may be affected significantly: we assume this does not hold.

Equation (19) is the main result of the paper. For the models of section II, simply requiring a TeV gravitino mass, achieved by $\mathcal{V} \sim 10^{15}$, naturally generates the required suppression scale $\Lambda \sim 10^{14} \text{GeV}$ for neutrino masses. This result is appealing, as it ties the scale of neutrino masses to that of supersymmetry breaking in a consistent manner. It is also appealing that the scale of supersymmetry breaking is naturally hierarchically small, as the moduli stabilisation mechanism of [6] generates exponentially large volumes.

We can condense the arguments of this section into a single formula relating neutrino masses, Higgs parameters, supersymmetry breaking and the gauge coupling

constants. Up to $\mathcal{O}(1)$ factors in $Z_{\alpha\beta\gamma\delta}$, we find

$$m_\nu \simeq \frac{v^2 \sin^2 \beta (\alpha_{SM}(m_s))^{2/3}}{2M_P^{2/3} (m_{3/2})^{1/3}}. \quad (20)$$

III. ON INTEGRATING OUT HEAVY FIELDS

A surprising result is that although the string scale is $m_s \sim 10^{11} \text{GeV}$, the suppression scale is $\Lambda \sim 10^{14} \text{GeV}$. With an intermediate string scale 10^{11}GeV there exist heavy states integrated out in producing the low energy theory. These include excited string states ($m_s \sim 10^{11} \text{GeV}$) and Kaluza-Klein states ($m_{KK} \sim 10^9 \text{GeV}$). Naively such states generate dimension-five operators suppressed, in contrast to (19), only by m_s or m_{KK} . However in fact the suppression scale (19) arises independently of the mass scale that has been integrated out.

To show this, we imagine including in the Lagrangian an explicitly massive field Φ with mass $m_\Phi \sim \mathcal{V}^{-\alpha} M_P$. For stringy states $\alpha = 1/2$ and for KK states $\alpha = 2/3$. The superpotential and Kähler potential are

$$W = H_2 L \Phi + M_P \Phi^2, \quad (21)$$

$$K = \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}} H_2 \bar{H}_2 + \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}} L \bar{L} + \mathcal{K}_{\Phi\bar{\Phi}} \Phi \bar{\Phi}. \quad (22)$$

Then $m_\Phi^2 = e^{\hat{K}} \frac{M_P^2}{\mathcal{K}_{\Phi\bar{\Phi}}^2}$ and using (16) $\mathcal{K}_{\Phi\bar{\Phi}} \sim \mathcal{V}^{\alpha-1}$. Neither m_s , m_{KK} nor any function of $\mathcal{V}$ may enter (21) - as with (10), such a scale would introduce a non-holomorphic dependence on the moduli.

The canonically normalised $H_2 L \Phi$ operator is

$$\hat{Y}_{H_2 L \Phi} = e^{\hat{K}/2} \frac{Y_{H_2 L \Phi}}{\sqrt{K_{\Phi\bar{\Phi}} K_{L\bar{L}} K_{H_2 \bar{H}_2}}} = \mathcal{V}^{1/6-\alpha/2} Y_{H_2 L \Phi}.$$

The $H_2 L H_2 L$ operator arises through integrating out Φ to generate a fermion-fermion-scalar-scalar interaction, with an intermediate fermionic $\tilde{\Phi}$. With all fields canonically normalised, the fermionic propagator is $\sim 1/m_\Phi$ and the resulting dimensionful interaction strength is $\mathcal{V}^{1/6-\alpha/2} \times \frac{\mathcal{V}^\alpha}{M_P} \times \mathcal{V}^{1/6-\alpha/2} = \frac{\mathcal{V}^{1/3}}{M_P}$, independent of the mass scale integrated out. This is exactly the suppression scale found in (18). There is a sense in which stringy or KK states play the role of right-handed neutrinos, although by their nature such stringy fields cannot form part of the low-energy theory.

This result is striking independent of its relevance for neutrino masses: the suppression scale of a dimension-five operator can be entirely independent of the masses of the heavy states integrated out to produce the operator.

The above arguments apply to explicitly massive states such as stringy, Kaluza-Klein or winding states. There may also exist dangerous heavy states with masses attained in the field theory (e.g. through a Higgs mechanism): an example of such a state is a field-theoretic right-handed neutrino with a mass of 10^5GeV . The existence of such states is model-dependent and we simply assume these are absent.

IV. ON STRING REALISATIONS

In most of the original (Type II) semi-realistic string Standard Model constructions, starting with [20], lepton number is perturbatively conserved, forbidding neutrino Majorana mass terms. However these can be generated by either GUT-like mechanisms (e.g. [19]) or nonperturbative brane instanton terms like $e^{-T}\nu_R\nu_R$ [21, 22]. $U(1)_L$ is anomalous; the field T is charged and transforms as $T \rightarrow T - 2iQ$ while the neutrinos transform as $\nu_R \rightarrow e^{iQ}\nu_R$, making this term gauge invariant. Such an approach cannot be relevant here: both because the instanton amplitudes $e^{-T} \sim \frac{1}{\mathcal{V}}$ are too small and because the string scale $m_s \sim 10^{11}\text{GeV}$ restricts the maximal field theory mass of the right-handed neutrino.

We have instead assumed the H_2H_2LL operator to be allowed - i.e. lepton number is not a perturbative symmetry - and computed its suppression scale. The original work of [20] notwithstanding, recent approaches show that there is no principled objection to SM-like constructions in string theory violating lepton number [23].

Given the absence of a conventional field theory seesaw in large volume compactifications, we find the attractive accordance of scales found here notable. Following our arguments, for any large volume stringy construction (in principle but not necessarily as in section II) in which the $HHLL$ operator enters the superpotential with an $\mathcal{O}(1)$ coefficient, the simple requirement of TeV scale supersymmetry guarantees the correct scale for

neutrino masses. We find this strong connection between TeV supersymmetry and neutrino masses far from trivial, and regard it as motivation for the development of global string models in which all the details take place.

V. CONCLUSIONS

This note argues that the mysterious energy scale $\Lambda \sim 10^{14}\text{GeV}$ of Majorana neutrino masses has a clean origin, associated with geometric features of a large class of compactification manifolds. We require (i) perturbative lepton number violation; (ii) a geometry consisting of a large six dimensional compactification manifold of volume $\mathcal{V} \sim 10^{15}$ in string units (as required for TeV-scale supersymmetry), with the Standard Model on a small four-dimensional submanifold; (iii) a superpotential independent of $\mathcal{V}$, as holds in string compactifications.

This suppression scale is independent of the masses of heavy states that are integrated out. Given these assumptions one is led naturally to Majorana neutrino masses of order $10^{-2}\text{eV} - 10^{-1}\text{eV}$.

Acknowledgments

We thank C. Burgess, L. Ibáñez and F. Quevedo for conversations. JC thanks the Perimeter Institute and the ICTP, Trieste for hospitality while this work was ongoing.

-
- [1] R. N. Mohapatra *et al.*, arXiv:hep-ph/0510213.
 - [2] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
 - [3] N. Arkani-Hamed and Y. Grossman, Phys. Lett. B **459**, 179 (1999) [arXiv:hep-ph/9806223].
 - [4] Y. Grossman and M. Neubert, Phys. Lett. B **474**, 361 (2000) [arXiv:hep-ph/9912408].
 - [5] T. Gherghetta, Phys. Rev. Lett. **92**, 161601 (2004) [arXiv:hep-ph/0312392].
 - [6] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, JHEP **0503**, 007 (2005) [arXiv:hep-th/0502058].
 - [7] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429** (1998) 263 [arXiv:hep-ph/9803315].
 - [8] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436** (1998) 257 [arXiv:hep-ph/9804398].
 - [9] K. R. Dienes, E. Dudas and T. Gherghetta, Nucl. Phys. B **557** (1999) 25 [arXiv:hep-ph/9811428].
 - [10] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and J. March-Russell, Phys. Rev. D **65** (2002) 024032 [arXiv:hep-ph/9811448].
 - [11] K. Benakli, Phys. Rev. D **60** (1999) 104002 [arXiv:hep-ph/9809582].
 - [12] C. P. Burgess, L. E. Ibanez and F. Quevedo, Phys. Lett. B **447** (1999) 257 [arXiv:hep-ph/9810535].
 - [13] J. P. Conlon, F. Quevedo and K. Suruliz, JHEP **0508** (2005) 007 [arXiv:hep-th/0505076].
 - [14] J. P. Conlon, D. Cremades and F. Quevedo, arXiv:hep-th/0609180.
 - [15] J. P. Conlon, S. S. Abdussalam, F. Quevedo and K. Suruliz, arXiv:hep-th/0610129.
 - [16] S. B. Giddings, S. Kachru and J. Polchinski, Phys. Rev. D **66**, 106006 (2002) [arXiv:hep-th/0105097].
 - [17] K. Becker, M. Becker, M. Haack and J. Louis, JHEP **0206**, 060 (2002) [arXiv:hep-th/0204254].
 - [18] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0301240].
 - [19] M. Cvetič and P. Langacker, arXiv:hep-th/0607238.
 - [20] L. E. Ibanez, F. Marchesano and R. Rabadan, JHEP **0111**, 002 (2001) [arXiv:hep-th/0105155].
 - [21] R. Blumenhagen, M. Cvetič and T. Weigand, arXiv:hep-th/0609191.
 - [22] L. E. Ibanez and A. M. Uranga, arXiv:hep-th/0609213.
 - [23] P. Anastasopoulos, T. P. T. Dijkstra, E. Kiritsis and A. N. Schellekens, arXiv:hep-th/0605226.
 - [24] $\mathcal{V}$ is the dimensionless volume, $\mathcal{V} = \text{Vol}/l_s^6$, with $l_s = 2\pi\sqrt{\alpha'}$. Ω is the (3,0)-form of the Calabi-Yau $\mathcal{M}$, S and T are dilaton and Kähler moduli superfields, g_s is the string coupling and G_3 the 3-form flux. A_i and a_i are $\mathcal{O}(1)$ numbers and $\xi \sim \chi(M)$ accounts for α' corrections.
 - [25] This assumes the superpotential Yukawas are independent of $\mathcal{V}$, which holds for all stringy embeddings [14].